



Contribution ID: 14

Type: **not specified**

## LHC-spin: a polarized internal target for the LHC

*Thursday, 26 September 2019 11:00 (20 minutes)*

We discuss the application of an open storage cell as gas target for a proposed LHC fixed-target experiment LHC-spin. The target provides a high areal density at minimum gas input, which may be polarized 1H, 2H, or 3He gas or heavy inert gases in a wide mass range. For the study of single-spin asymmetries in pp interaction, luminosities of nearly  $10^{33}/\text{cm}^2 \text{ s}$  can be produced with existing techniques.

### Summary

We discuss the application of an open storage cell as gas target for a proposed LHC fixed-target experiment LHC-spin. The target provides a high areal density at minimum gas input, which may be polarized 1H, 2H, or 3He gas or heavy inert gases in a wide mass range. For the study of single-spin asymmetries in pp interaction, luminosities of nearly  $10^{33}/\text{cm}^2 \text{ s}$  can be produced with existing techniques.

**Primary author:** LENISA, Paolo (University of Ferrara and INFN - Ferrara (Italy))

**Presenter:** LENISA, Paolo (University of Ferrara and INFN - Ferrara (Italy))

**Session Classification:** Polarized Gas Targets

**Track Classification:** Polarized Gas Targets